

# Turfgrass Consumptive Use Values for the Phoenix Area

## Introduction


Consumptive use (CU) curves that provide average rates of turfgrass evapotranspiration ( $ET_T$ ) are widely used by irrigation professionals for design and management of turfgrass irrigation systems. For approximately 35 years, the bermudagrass lawn CU curve developed by the United States Department of Agriculture (Erie et al., 1965; Erie et al., 1982) has served as the lone published CU curve for turfgrass in Arizona. While the USDA CU curve has proven useful to the turf industry, turf professionals do question whether  $ET_T$  values obtained from the curve are relevant to turf systems commonly used in Arizona today. The USDA curve was developed for the summer turf season using a low-maintenance common bermudagrass mowed to a height of 3.8 cm (1.5") every four weeks, and watered every two weeks using flood irrigation (Garrot and Mancino, 1994). A relevant turf system today consists of hybrid bermudagrass maintained at a height of ~2 cm (0.75") and watered at frequent intervals using sprinkler irrigation. The practice of overseeding with ryegrass in the fall to maintain green cover in winter is also common today. The USDA CU curve does not address the issue of overseeding and provides no information on  $ET_T$  for the period mid-October through mid-April.

A number of research studies have been completed in recent years to quantify the water requirements of turfgrass grown in the low desert regions of Arizona (Brown et al., 1996; Brown et al., 2001). Several studies had as their primary objective the development of crop coefficients (Kcs) that convert reference evapotranspiration ( $ET_o$ ) data computed from meteorological data (from weather stations) into estimates of  $ET_T$  (Brown et al., 2001). In this report, we apply Kcs developed from these studies to long-term records of  $ET_o$  to provide updated CU information for turfgrass grown in the Phoenix metropolitan area.

## Methods

Estimates of  $ET_T$  were computed on a daily basis for the period 1987 through 2000 by applying turfgrass Kcs to the historical record of reference evapotranspiration ( $ET_o$ ) available for the Phoenix area from the Arizona Meteorological Network (AZMET). Specific

## PHOENIX



— High Quality — Accept. Quality

Figure 1. Consumptive use curves for high and acceptable quality turf grown in the Phoenix area.

data used in this analysis were obtained from two Phoenix area AZMET stations located on golf courses: Phoenix Greenway (Cave Creek Golf Course) and Phoenix Encanto (Encanto Golf Course). The mathematical procedure used to produce the  $ET_T$  estimates involved multiplying the appropriate crop coefficient (Kcs) by  $ET_o$ :

$$ET_T = Kcs \times ET_o$$

4/2003

AZ1314

### Turf Irrigation Management Series: V

THE UNIVERSITY OF ARIZONA  
COLLEGE OF AGRICULTURE AND LIFE SCIENCES  
TUCSON, ARIZONA 85721

Paul Brown  
Biometeorology Specialist

*This information has been reviewed by university faculty.*

[cals.arizona.edu/pubs/water/az1314.pdf](http://cals.arizona.edu/pubs/water/az1314.pdf)

Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, James A. Christenson, Director, Cooperative Extension, College of Agriculture & Life Sciences, The University of Arizona.

The University of Arizona is an equal opportunity, affirmative action institution. The University does not discriminate on the basis of race, color, religion, sex, national origin, age, disability, veteran status, or sexual orientation in its programs and activities.

The Kcs used to estimate  $ET_T$  were developed for a common desert turf system consisting of Tifway bermudagrass in summer and overseeded ryegrass in winter. Other assumptions implicit in the use of the Kcs employed include frequent irrigation with sprinklers, mowing heights ranging from 0.625-1.0" in summer and 0.875-1.25" in winter, and two levels of turf quality defined as high and acceptable. High quality turf areas would include high profile sports turf (playing fields and golf course fairways) and areas where turf appearance is very important. These areas generally receive high levels of fertilization and maintenance. Acceptable quality turf would be suitable for lawn or park environments where traffic is low, rapid regrowth is not required and fertilization levels are relatively low.

Crop coefficients appropriate for high quality turf were based on the research results of Brown et al. (2001) and change monthly. Crop coefficients for acceptable quality turf were derived by subtracting 0.1 from the high quality Kcs (Brown et al., 1996).

The resulting 14 years of daily  $ET_T$  data from the two AZMET sites were first averaged by day of the year to produce an average annual  $ET_T$  data set for each location. These location specific  $ET_T$  values were in turn averaged to produce a Phoenix-area annual

$ET_T$  data set. This Phoenix-area daily  $ET_T$  data set was then summarized into weekly, monthly, and annual totals of  $ET_T$ . Consumptive use curves were developed for high and acceptable quality turf from the summarized data sets.

## Results

Annual CU curves for high and acceptable quality turfgrass grown in Phoenix area are presented in Figure 1. Turfgrass ET varies nearly 5-fold over the course of the year, reflecting the annual fluctuation in atmospheric evaporative demand. The ET of high quality turf ranges from a low of ~0.05"/day in December to ~0.25"/day in June. Evapotranspiration from acceptable quality turf runs about 15% below that of high quality turf and ranges from ~0.04"/day in December to ~0.22"/day in June.

Weekly as opposed to daily values of  $ET_T$  may prove more useful when managing irrigation, especially if irrigation is not being applied each day. Table 1 provides weekly totals of  $ET_T$  for high and acceptable quality turfgrass grown in the Phoenix area. Evapotranspiration from high quality turf ranges from 0.31" in the first week of January to 1.83" in the first week of July. The range in weekly  $ET_T$  for acceptable quality turf

Table 1. Weekly consumptive use ( $ET_T$ ) in inches for high and acceptable quality turf grown in the Phoenix area

Week Ending	Turf Quality										
	High	Acc.									
Jan 7	0.31"	0.26"	Apr 8	1.23"	1.07"	Jul 8	1.83"	1.59"	Oct 7	1.13"	0.98"
Jan 14	0.37	0.31	Apr 15	1.40	1.22	Jul 15	1.73	1.51	Oct 14	1.04	0.90
Jan 21	0.41	0.35	Apr 22	1.43	1.24	Jul 22	1.71	1.49	Oct 21	0.92	0.80
Jan 28	0.43	0.37	Apr 29	1.53	1.33	Jul 29	1.71	1.49	Oct 28	0.75	0.65
Feb 4	0.52	0.44	May 6	1.59	1.38	Aug 5	1.78	1.56	Nov 4	0.70	0.60
Feb 11	0.53	0.45	May 13	1.63	1.41	Aug 12	1.72	1.52	Nov 11	0.62	0.54
Feb 18	0.56	0.48	May 20	1.65	1.43	Aug 19	1.68	1.48	Nov 18	0.56	0.49
Feb 25	0.67	0.57	May 27	1.67	1.45	Aug 26	1.58	1.39	Nov 25	0.53	0.46
Mar 4	0.71	0.61	Jun 3	1.71	1.48	Sep 2	1.42	1.24	Dec 2	0.48	0.41
Mar 11	0.83	0.72	Jun 10	1.68	1.45	Sep 9	1.37	1.20	Dec 9	0.39	0.34
Mar 18	0.90	0.78	Jun 17	1.73	1.49	Sep 16	1.33	1.16	Dec 16	0.38	0.33
Mar 25	1.03	0.89	Jun 24	1.72	1.48	Sep 23	1.24	1.08	Dec 23	0.33	0.28
Apr 1	0.93	0.81	Jul 1	1.81	1.56	Sep 30	1.22	1.06	Dec 31*	0.38	0.33

\* Water use for the week ending December 31 represents an 8-day total for the period December 24-31.

ranges from 0.26" in early January to 1.59" in early July.

Monthly values of  $ET_T$  are useful when planning irrigation budgets for a year. Table 2 presents monthly  $ET_T$  for high and acceptable quality turfgrass for the Phoenix area. Monthly  $ET_T$  for high quality turf ranges from 1.6" in December to 7.7" in July. For acceptable quality turf,  $ET_T$  ranges from 1.4" in December to 6.7" in July. The last column in Table 2 presents the percentage of annual  $ET_T$  occurring in each month. These monthly percentages clearly show that the bulk of the annual water use occurs during the summer months. For example,  $ET_T$  in July accounts for 13.4% of total annual  $ET_T$ . In contrast, total  $ET_T$  from December through February (inclusive) represents just 9.9% of annual  $ET_T$  — substantially less than  $ET_T$  for July.

Annual CU of high and acceptable quality turf is summarized at the bottom of Table 2. Consumptive use of high quality turf totals ~57.7" or 4.8' per year while the CU of acceptable quality turf approaches ~49.9" or 4.16' per year.

Table 2. Monthly and annual consumptive use ( $ET_T$ ) in inches for high and acceptable quality turf grown in the Phoenix area.

Month	Turf Water Use		
	Turf Quality		% of Annual Use
	High	Acc.	
January	1.7"	1.4"	3.0
February	2.4	2.0	4.1
March	4.0	3.4	6.9
April	5.9	5.2	10.3
May	7.3	6.3	12.7
June	7.4	6.4	12.8
July	7.7	6.7	13.4
August	7.3	6.4	12.7
September	5.6	4.8	9.6
October	4.2	3.6	7.2
November	2.5	2.1	4.3
December	1.6	1.4	2.8
Total	57.5"	49.9"	

## Concluding Remarks

The CU data presented in this report represent long-term average rates of  $ET_T$  and should prove useful to individuals involved in the design and management of turf irrigation systems. It is important to realize that the results presented in this report represent raw  $ET_T$  data that have not been adjusted for precipitation or irrigation system performance. To use this CU information to determine the amount of water required for irrigation, one must first subtract the amount of effective precipitation (precipitation not lost to deep percolation and runoff) to determine the net water requirement for any period. Precipitation in the Phoenix area averages ~7" or 0.6' per year and should reduce irrigation water requirements to some degree.

The final step in determining irrigation water requirements involves making adjustments to: 1) account for system nonuniformity and 2) ensure leaching is sufficient to maintain soil salinity at acceptable levels. Adjustments for nonuniformity and salinity management increase the amount of irrigation water required and vary dramatically with location due to differences in irrigation design, topography, local weather conditions, and water quality. A discussion of these adjustments is beyond the scope of this publication and will be discussed in a subsequent report in the Turf Irrigation Management Series.

## References

- Brown, P.W., J. Gilbert, and D.M. Kopec. 1996. Turfgrass irrigation scheduling using weather based estimates of evapotranspiration for high and low traffic turfs. Final Report to: AZ Dept. Water Resources. Contract No. CA94TU103-00.
- Brown, P.W., C.F. Mancino, T.L. Thompson, M.H. Young, P.J. Wierenga, and D.M. Kopec. 2001. Penman Monteith crop coefficients for desert turf systems. *Crop Sci.* 41: 1197-1206.
- Erie, L.J., O.F. French, and K. Harris. 1965. Consumptive use of water by crops in Arizona. *Tech. Bul.* 169. Agricultural Exp. Stn., Univ. of Arizona. 46p.
- Erie, L.J., O.F. French, D.A. Bucks, and K. Harris. 1982. Consumptive use of water by major crops in the southwestern United States. *USDA Cons. Res. Rpt.* 29. Washington, DC.
- Garrot, D.J., and C.F. Mancino. 1994. Consumptive water use of three intensively managed bermudagrasses growing under arid conditions. *Crop Sci.* 34: 215-221.

---

*Any products, services, or organizations that are mentioned, shown, or indirectly implied in this publication do not imply endorsement by The University of Arizona.*